
The WAGER, Vol. 1(6) - Pathological gambling among methadone patients

February 6, 1996

Research suggests that substance-abusing populations have higher rates of problem and pathological gambling than the general adult population. A recent study examined the gambling behavior of a random sample (n = 117) of respondents enrolled in a large methadone maintenance treatment program. The study found that methadone patients had higher rates of problem (15%) and probable pathological (16%) gambling than the general adult population. Although this study was based on a small sample from a single treatment program, other studies of substance-abusing patients provide support for these findings. For example, previous studies have found pathological gambling rates of 9% among patients seeking substance abuse treatment (Lesieur, Blume & Zoppa, 1986), 11% among patients with a secondary diagnosis of substance abuse (Lesieur & Blume, 1990), and 14% among young substance abusers in a therapeutic community program (Lesieur & Heinemann, 1988).



Sources:

Lesieur, H.R. & Blume, S.B. (1990). Characteristics of pathological gamblers identified among patients on a psychiatric admissions service. Hospital and

Community Psychiatry, 41, 1009-1012;

Lesieur, H.R., Blume, S.B., & Zoppa, R.M. (1986). Alcoholism, drug abuse, and gambling. Alcoholism: Clinical and Experimental Research, 10, 33-38

Lesieur, H.R. & Heineman, M. (1988). Pathological gambling among youthful multiple substance abusers in a therapeutic community. British Journal of Addiction, 83, 765-771

Spunt, B., Lesieur, H., Hunt, D., & Cahill, L. (1995). Gambling among methadone patients. International Journal of the Addictions, 30, 929-962.

This public education project is funded, in part, by the Massachusetts Department of Public Health.

This fax may be copied without permission. Please cite The WAGER as the source.

For more information contact the Massachusetts Council on Compulsive Gambling, 190 High Street, Suite 6, Boston, MA 02110.