

ASHES, Vol. 15(10) - Can social media be the answer to quitting smoking? The effects of a social media campaign on young adult smoking cessation

September 24, 2019

Social media platforms are [extremely popular in the United States](#) and worldwide. While social media has been implicated in [increasing rates of depression](#) and [excessive internet use](#) among young adults, it might also help people manage addictive behaviors. This week, as part of our [Special Series on Addiction and Social Media](#), ASHES reviews [a study by Neill Baskerville and colleagues](#) that examined how a social media campaign influenced the chance of successfully quitting smoking.

What was the research question?

Is a social media-based smoking cessation program more effective at helping young adults quit smoking than a conventional helpline smoking cessation program?

What did the researchers do?

The researchers recruited 238 young adult smokers from Canada to take a survey and participate in a program to quit smoking. They exposed¹ 102 participants to a social media-based smoking cessation program and 136 participants to a helpline cessation program. Using logistic regression models and controlling for several potential confounders², they analyzed how the two types of smoking cessation programs influenced: 1) attempting to quit smoking³, 2) quitting smoking for 7 consecutive days; and 3) quitting smoking for 30 consecutive days.

What did they find?

People who were exposed to the social media campaign were more successful at quitting smoking than people who used the smoking helpline. The social media group had 2.05 [greater odds](#) of making a quit attempt than the smoking helpline

group (see figure). Participants in the social media group also had 2.14 greater odds of remaining smoke-free for 30 days.



Figure. Odds of making a quit attempt and successfully quitting smoking for 7 and 30 days for two types of smoking cessation programs. Click image to enlarge.

Why do these findings matter?

While much of the focus of research on social media has examined potential adverse effects, this study shows that using social media to help people quit smoking might work better than at least one conventional method (i.e., a smoking helpline). These findings are important because this technological approach to reducing smoking could be extended to other types of addictive behaviors. Other studies have found social media campaigns to be successful in [reducing stigma related to mental health issues](#) and for [recruiting high-risk substance users for research](#), suggesting that social media has promise for enhancing addiction treatment and research.

Every study has limitations. What are the limitations in this study?

As the researchers used [self-reported](#) survey data on smoking cessation, this measure may not accurately measure actual smoking behavior. The study sample was relatively small and was not nationally [representative](#), so these results are not likely to [generalize](#) to all smokers.

For more information:

[SmokeFree](#) offers tools and tips about quitting and maintaining abstinence from smoking. The [Centers for Disease Control and Prevention](#) provides research and tips about cigarettes and how to quit. For evidence-based information about the health effects of smoking cigarettes, read [this page hosted by the American Cancer Society](#). For additional tools, please visit the BASIS [Addiction Resources](#) page.

— Eric R. Louderback, Ph.D.

What do you think? Please use the comment link below to provide feedback on this article.

1. This study used a quasi-experimental design, without [randomly assigning participants](#) to each type of smoking cessation program. Although the authors compared the demographics of the groups and adjusted for group differences, their results are likely not as accurate as a [controlled experimental design](#).

2. To ensure that other potential variables that could be associated with quitting smoking did not influence their results, the authors controlled for education level, frequency of cigarette use, and ethnicity based on a survey administered to participants at the beginning of the study.

3. The authors measured “making a quit attempt” as successfully stopping smoking for 24 or more consecutive hours during the study period.