

# The DRAM, Vol. 16(5) - Do state tax rates on alcohol and responsible drinking policies reduce risky alcohol use?

April 28, 2020

Drinking in excess is associated with numerous consequences [including psychological and physical health problems, traffic fatalities and alcohol use disorder](#). In the U.S., [rates of binge drinking](#) (i.e., more than 4 drinks per occasion for women or more than 5 drinks per occasion for men) are persistently high, despite widespread public health initiatives. Despite policies<sup>1</sup> designed to encourage responsible drinking behaviors, little is known about their impact on risky alcohol use. In this week's DRAM, we review [a study by Brian Fairman and colleagues](#) that examined how 19 state-level alcohol policies in the United States might be associated with binge drinking patterns in adolescence and young adulthood.

## **What were the research questions?**

What are the effects of state-level alcohol taxes, responsible drinking policies and local liquor store availability on trajectories of drinking among young adults?

## **What did the researchers do?**

The researchers gathered three types of information: survey responses from young people from 22 U.S. states for every year over a five-year period, starting in 2010 when they were in 10th grade; descriptions of their states' alcohol-related policies<sup>2</sup> and beer excise tax rates; and the number of liquor stores within 1 kilometer of each respondent's home. Next, they used [group-based trajectory modeling](#) to determine: (1) how often each respondent engaged in binge drinking during each year between 2010 and 2015, and (2) if there were differences in these drinking patterns over time based on a participant's state-level alcohol policies and proximity to liquor stores in their community.

## **What did they find?**

The authors found that there were five distinct patterns of binge drinking over the

five year period: **low-risk drinking** (i.e., not engaging in binge drinking), **escalating** (i.e., increasing binge drinking frequency), **late-onset** (i.e., increasing binge drinking in later years), **chronic** (i.e., consistent high levels of binge drinking) and **decreasing** (i.e., a reduction in binge drinking). The largest group of respondents were in the low-risk category (32.9%), followed by the escalating category (26.1%) and the chronic category (15.1%). Respondents who lived in states with weaker alcohol policies were more likely to be in the escalating group, while respondents who lived in states with higher beer excise taxes were less likely to be in the escalating group. Distance to liquor stores in one's neighborhood did not significantly predict binge-drinking frequency.




Figure. Figure shows beer excise tax rates for 2020 (dollars per gallon of alcohol) as well as underaged alcohol possession and driving license revocation laws for 2020 in the contiguous United States. Click figure to enlarge.

Sources: <https://www.taxpolicycenter.org/statistics/state-alcohol-excise-taxes>; Taxes:  
<https://alcoholpolicy.niaaa.nih.gov/apis-policy-topics/uselose-driving-privileges/44/maps-and-charts#page-content>. Use/Lose Laws:

### Why do these findings matter?

The results are important for developing public health interventions to reduce binge drinking and other types of risky alcohol use. Policies and excise taxes appear to be more important for predicting drinking behaviors than liquor store availability, so states should consider these findings when developing policies. For

example, [enacting zero tolerance laws for underage DUI](#) and [higher taxes on alcohol](#) might be effective in reducing risky drinking patterns among young adults. In addition to policy changes, successful interventions such as [social media-based programs to promote healthier alcohol use](#) and [the use of text messages to encourage responsible drinking](#) could potentially reduce unhealthy drinking patterns to an even greater degree.

### **Every study has limitations. What are the limitations in this study?**

While this study examined how policies are associated with drinking behavior, the relationship might not be [causal](#) because other [unmeasured confounding variables](#) could also have influenced binge drinking involvement. Also, using [self-reported data](#) may also reduce the [validity](#) of the findings, because respondents might not accurately report their drinking behaviors. Moreover, confusion about what defines “one standard drink” of alcohol might also lead to under- or overestimates of one’s actual alcohol intake and binge drinking frequency.

### **For more information:**

The National Institute for Alcohol Abuse and Alcoholism has [tips and resources](#) for people struggling with problem drinking. For drinking self-help tools, please visit The BASIS [Addiction Resources](#) page. For more information about Open Science practices, check out the [OSF website](#).

— Eric R. Louderback, Ph.D.

*What do you think? Please use the comment link below to provide feedback on this article.*

[1] Historically, states have implemented [several policies to reduce excessive drinking](#), especially among young people. These policies include a 21+ drinking age (passed in 1984), restrictions on public consumption of alcohol, and severe legal penalties for driving while intoxicated.

[2] These 19 state-level policies included: 1) motor vehicle regulations mandating license loss for underage violation of alcohol laws (i.e., use/lose laws), laws against 2) underage possession, 3) purchase, 4) consumption, 5) internal consumption, 6) possession of a fake identification and 7) creating fake identification, 8) whether Sunday alcohol sales are allowed, 9) if there is a state-controlled distribution system for alcoholic beverages, and 10 types of legal regulations for alcohol providers (e.g., bartenders and liquor stores).