

STASH, Vol. 16(8) - Have opioid overdose deaths been underreported?

August 4, 2020

Editor's Note: *Can you spare five minutes to tell us your thoughts on The BASIS? Your responses to this [short questionnaire](#) will help us improve this free resource and better meet your expectations. Thank you!*

The world has improved in recent years on many metrics- fewer people are [living in extreme poverty](#), [dying from air pollution](#), or [missing out on a basic education](#). One clear exception to this trend is opioid overdoses: The number of people who die each year from opioids [nearly doubled between 2007 and 2017](#). With the opioid epidemic as [the leading cause of death](#) in the United States for adults under 50 years old, the scope of its catastrophic effects seems difficult to underestimate. And yet, the U.S. probably undercounts the true number of fatal opioid overdoses in the U.S. Over 20% of overdose deaths [go unclassified each year](#), likely because coroners do not always have the expertise or resources to detect the presence of opioids in decedents. STASH reviews an study in which [Andrew Boslett and colleagues](#) compare various methods for estimating how many unclassified overdose deaths were caused by opioids.

What were the research questions?

How many unclassified overdoses deaths from 1999-2016 were caused by opioids? Which variables and statistical techniques should researchers use to determine whether opioids were involved in unclassified overdose deaths?

What did the researchers do?

Using 1999-2016 data from the (U.S.) Centers for Disease Control and Prevention's National Center for [Health Statistics Detailed Multiple Cause of Death](#) records, Boslett and his co-authors compared the relative importance of characteristics of the decedent (e.g., marital status), the presence of other factors that commonly contribute to opioid-related deaths (e.g., obesity), and characteristics of the county in which the death occurred (e.g., poverty rate) in predicting whether opioids were involved in an overdose death. The authors also compared two statistical methods for predicting opioid overdoses, [logistic](#)

[regression](#) and [random forests](#). Logistic regression performs best when there are linear relationships between predictors and outcome. For example, it could be that overdose deaths in poor counties are more often caused by opioids than overdose deaths in wealthy counties. Random forests excels at detecting complex relationships among the different pieces of information available to predict the outcome. For instance, perhaps dying by overdose in a poor area is only indicative of opioid use if the decedent was not married, was between the age of 20-30, and died during the weekend. After developing the algorithms using cases in which the substance involved in an overdose was specified, Boslett and his colleagues used the algorithms to make educated guesses about which unclassified overdose deaths involved opioids.

What did they find?

Including the presence of other common contributing causes of death (see Figure) and decedent characteristics in the models greatly improved the classification of overdose deaths as involving or not involving opioids. By contrast, county-level characteristics provided little added value. Random forests performed no better than logistic regression. The best algorithm found that 71.8%– or just under 100,000– of unclassified overdose deaths probably involved opioids.

Which contributing causes of death made opioid overdose more likely?

The presence of...	The absence of...
F111: Opioid abuse	S068: Intracranial injuries
F119: Opioid use, unspecified	S062: Diffuse traumatic brain injury
F112: Opioid dependence	I619: Nontraumatic intracerebral hemorrhage
R529: Pain, unspecified	E162: Hypoglycemia
M549: Dorsalgia, unspecified	F159: Other stimulant abuse, unspecified
R522: Other chronic pain	K720: Acute and subacute hepatic failure
M797: Fibromyalgia	F151: Other stimulant abuse
F192: Other psychoactive substance dependence	K729: Hepatic failure, unspecified
R098: Symptoms involving the circulatory or respiratory systems	I48: Atrial fibrillation and flutter
J180: Bronchopneumonia, unspecified organism	T751: Unspecified effects of drowning and nonfatal submersion

Figure. The [International Classification of Diseases \(ICD\)](#) codes that most strongly predict opioid involvement in overdose death. On the left are the ICD codes that are more common when opioids were involved in fatal overdoses; on the right are ICD codes that are usually not reported by coroners when opioids were involved in fatal overdoses. To learn more about a certain diagnosis, enter the alphanumeric code [here](#). Click image to enlarge.

Why do these findings matter?

Policymakers deciding how to use public funds should consider that the opioid epidemic may be 28% larger than official statistics indicate. To avoid underestimating opioid overdoses in the future, coroners need routine education and resources to detect opioid-related deaths. Finally, more complex algorithms may not be necessary for accurately estimating the true scope of the opioid epidemic.

Every study has limitations. What are the limitations of this study?

The authors only compared the performance of logistic regression on its own to random forests on its own. It may be that [other algorithms, or combinations of algorithms](#), would have had better performance than logistic regression. Also, if there are characteristics of unclassified overdose deaths that make them more difficult for coroners to classify even with adequate training and resources, then algorithms that were developed on classified deaths may be less valid for predicting opioid involvement in overdose deaths of unspecified origin.

For more information:

If you are worried that you or someone you know is experiencing addiction, the [SAMHSA National Helpline](#) is a free treatment and information service available 24/7. For more details about addiction, visit our [Addiction Resources](#) page.

— William McAuliffe, PhD

What do you think? Please use the comment link below to provide feedback on this article.