

ASHES, Vol. 22(10) - Is tobacco use among adolescents associated with poor dental health over time?

September 15, 2026

During adolescence, youth are having new experiences and adopting new habits, some of which may impact their oral health. Cavities and tooth decay are [common](#) oral health concerns among adolescents, which, if left [untreated](#), can lead to infection and even impact school performance and social development. Risky behaviors like [adolescent](#) tobacco and e-cigarette smoking could [contribute](#) to poor oral health, and the risk might be greater when these substances are [combined](#). However, longitudinal studies are needed to test the association between tobacco use and oral health outcomes among adolescents over time. This week, ASHES reviews a study by [Noor Abdulhay and colleagues](#) that did just that.

What was the research question?

Among adolescents, is tobacco use associated with developing oral health problems over time?

What did the researchers do?

The researchers analyzed data from 15,788 adolescents in the [longitudinal](#), six Wave, [Population Assessment of Tobacco and Health \(PATH\)](#) study. Waves 1-6 occurred between 2013-2021. Participants enter the study on a rolling basis – whichever wave they entered was considered their baseline. Via telephone and in-person interviews, participants reported past-month tobacco use and dental health concerns at baseline and subsequent waves. Participants were grouped by type of tobacco use (i.e., combustible tobacco products (CTP) only, e-cigarettes only, dual use, and no-use). Those who reported dental concerns at baseline were removed from the sample. [Weighted Cox regression](#) models were used to assess tobacco use and the risk of experiencing oral health problems, while also controlling for [covariates](#) like adolescent school level, parent marital status, parent education status, diabetes diagnosis, and tobacco access at home.

What did they find?

Adolescents who reported no tobacco use in the past month were less likely to develop oral health concerns over time, compared to those who used no tobacco

products. However, these heightened risks were limited to adolescents who used *e-cigarettes only* and those who engaged in *dual use* of e-cigarettes and CTPs (e.g., cigarettes/hookahs) – not those who only used CTPs. Some findings held true even when controlling for covariates. More specifically, adolescents who engaged in past-month *dual use* were more likely than those who did not use tobacco to develop oral health problems, after adjustment (see Figure).




Figure. Displays the [hazard odds](#) of reporting tobacco use status and developing a dental health problem over time. Odds ratios can be interpreted as having X times greater/lesser odds of reporting an outcome compared to a reference group. [Statistically significant](#) relationships are denoted by *. Click image to enlarge.

Why do these findings matter?

Any form of smoking can [negatively influence](#) oral health. For example, chemicals in e-cigarettes can [mimic](#) the effect that high-sugar foods and acidic drinks have on dental health. According to the findings in this study, dual use among adolescents is particularly harmful, emphasizing the need for greater public awareness of its impact. This could be achieved through [media campaigns](#), screening from [healthcare providers](#), [school outreach](#), and more. These efforts should not only target young people who use these products but also their [guardians](#) who are likely managing their dental care.

Every study has limitations. What are the limitations of this study?

Participants [self-reported](#) their tobacco use and dental concerns, making the results subject to [recall bias](#). Additionally, the dental health questionnaire did not distinguish between types of dental concerns, limiting the ability to determine the impact of types of tobacco use on specific dental health conditions. Finally, this dataset used covariates that were consistent across all waves. Therefore, researchers could not determine the impact of covariates like oral hygiene and socioeconomic factors that could influence dental health and access to care in different ways over time.

For more information:

Individuals who want to quit smoking or vaping may benefit from engaging with the [American Lung Association](#). Additional resources can be found at the BASIS [Addiction Resources](#) page.

— Nakita Sconsoni, MSW

Want CE credit for reading BASIS articles? [Click here](#) to visit our Courses Website and access our free online courses.
